



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

botanists a number of very essential facts. The present "Notes" deal with four specimens of *Lepidodendron harcourtii* and *Halonias regularis*, which the author regards as specifically identical; and with a new genus of Cycadofilices to which he assigns the name of *Megaloxylon*, as represented by a single species, *M. scottii*.

D. P. P.

Matonia pectinata.¹—In an important contribution to the *Philosophical Transactions of the Royal Society*, which gives evidence of the most searching care, Mr. Seward selects *Matonia* as representing an isolated type of ferns requiring further examination anatomically, in order to determine its relations to the Cyatheaceæ and Gleicheniaceæ, toward which it has been customary to assign it an intermediate position. Mr. Seward concludes that while the genus must hold an independent position among ferns, its affinities are most nearly with those of the Cyatheaceæ. His studies of existing species are designed to have a direct bearing upon the geological history of the Matonineæ, and he therefore brings into comparison *Laccopteris* and *Matonidium* in particular; from all of which he concludes that *Matonia* represents the survival of a family of ferns now confined to a few localities in Borneo and the Malay Peninsula, and represented by two living species, which in Mesozoic time had a wide geographical range, being especially abundant in the European area during the Jurassic and Lower Cretaceous times. The apparent absence of the Matonineæ from Tertiary formations suggests that these forms reached their maximum development in the Mesozoic, and that toward the close of the Cretaceous a decided restriction in geographical range had developed.

D. P. P.

Medullosa anglica.²—Dr. Scott's important contributions to our knowledge of the Cycadofilices have been recently enlarged by the description of a new species of *Medullosa* to which he assigns the name of *M. anglica*. He points to the fact that several genera such as *Nœggerathia*, *Medullosa*, *Cladoxylon*, *Lyginodendron*, *Heterangium*, and *Protopytis* are now to be grouped under Potonie's con-

¹ Seward, A. C. The Structure and Affinities of *Matonia pectinata*, R. Br., with Notes on the Geological History of the Matonineæ, *Phil. Trans. Roy. Soc.*, Ser. B, vol. cxc (1899), pp. 171.

² Scott, D. H. On the Structure and Affinities of Fossil Plants from the Palæozoic Rocks. III. *Medullosa anglica*, a new representative of the Cycadofilices, *Phil. Trans. Roy. Soc.*, Ser. B, vol. cxc (1899), pp. 81.